


Keeling Curve

“Keeling” redirects here. For other uses, see [Keeling \(disambiguation\)](#).

The **Keeling Curve** is a graph which plots the ongoing



The Keeling Curve, December 2014, Scripps Institution of Oceanography, UC San Diego

ing change in concentration of carbon dioxide in Earth’s atmosphere since 1958. It is based on continuous measurements taken at the Mauna Loa Observatory in Hawaii that began under the supervision of Charles David Keeling. Keeling’s measurements showed the first significant evidence of rapidly increasing carbon dioxide levels in the atmosphere. Many scientists credit Keeling’s graph with first bringing the world’s attention to the current increase of carbon dioxide in the atmosphere.^[1]

Charles David Keeling, of Scripps Institution of Oceanography at UC San Diego, was the first person to make frequent regular measurements of the atmospheric carbon dioxide (CO₂) concentration, taking readings at the South Pole and in Hawaii from 1958 onwards.^[2]

Prior to Keeling, the concentration of carbon dioxide in the atmosphere was thought to be affected by constant variability. Keeling had perfected the measurement techniques and observed “strong diurnal behavior with steady values of about 310 ppm in the afternoon” at three locations: (Big Sur near Monterey, the rain forests of Olympic Peninsula and high mountain forests in Arizona).^[3] By measuring the ratio of two isotopes of carbon, Keeling attributed the diurnal change to respiration from local plants and soils, with afternoon values representative of the “free atmosphere”. By 1960, Keeling and his group had determined that the measurement records from California, Antarctica, and Hawaii were long enough to see not just the diurnal and seasonal variations, but also a year-on-year increase that roughly matched the amount of fossil fuels burned per year. In the article that made him famous, Keeling observed, “at the South Pole the ob-

served rate of increase is nearly that to be expected from the combustion of fossil fuel”.^[4]

1 Mauna Loa measurements



The Mauna Loa Observatory

Due to funding cuts in the mid-1960s, Keeling was forced to abandon continuous monitoring efforts at the South Pole, but he scraped together enough money to maintain operations at Mauna Loa, which have continued to the present day^[5] alongside the monitoring program by NOAA.^[6]

The measurements collected at Mauna Loa show a steady increase in mean atmospheric CO₂ concentration from about 315 parts per million by volume (ppmv) in 1958 to 401 ppmv as of April 2014.^{[7][8]} This increase in atmospheric CO₂ is considered to be largely due to the combustion of fossil fuels, and has been accelerating in recent years. Since carbon dioxide is a greenhouse gas, this has significant implications for global warming. Measurements of carbon dioxide concentration in ancient air bubbles trapped in polar ice cores show that mean atmospheric CO₂ concentration has historically been between 275 and 285 ppmv during the Holocene epoch (9,000 BCE onwards), but started rising sharply at the beginning of the nineteenth century.^[9]

Keeling and collaborators made measurements on the incoming ocean breeze and above the thermal inversion layer to minimize local contamination from volcanic vents. In addition, the data are normalized to negate any influence from local contamination.^[10] Measurements at many other isolated sites have confirmed the long-term trend shown by the Keeling Curve,^[11] though no sites have a record as long as Mauna Loa.^[12]

The Keeling Curve also shows a cyclic variation of about 5 ppmv in each year corresponding to the seasonal change in uptake of CO₂ by the world’s land vegetation. Most of

this vegetation is in the Northern hemisphere, since this is where most of the land is located. From a maximum in May, the level decreases during the northern spring and summer as new plant growth takes carbon dioxide out of the atmosphere through photosynthesis. After reaching a minimum in October, the level rises again in the northern fall and winter as plants and leaves die off and decay, releasing the gas back into the atmosphere.^[13]

Due in part to the significance of Keeling's findings,^[5] the NOAA began monitoring CO₂ levels worldwide in the 1970s.^[6] Today, CO₂ levels are monitored at about 100 sites around the globe.^[1]

Carbon dioxide measurements at the Mauna Loa observatory in Hawaii are made with a type of infrared spectrophotometer called a capnograph by, John Tyndall, its inventor in 1864, but now known as a nondispersive infrared sensor.^[14] Currently (May, 2013) several laser-based sensors are being added to run concurrently with the IR spectrophotometer at Scripps, while NOAA measurements at Mauna Loa use nondispersive infrared sensor tightly calibrated using WMO standards as the main reference for the curve.^[15] Multiple other sensors and technologies are also used at Mauna Loa to augment these measurements.

Keeling died in 2005. Supervision of the measuring project was taken over by his son, Ralph Keeling, a professor of geochemistry at Scripps Oceanography.^[16]

When C.D. Keeling published his 1960 paper, there did not appear to be any evidence of oceanic absorption. Keeling conjectured that this might be attributable to fluctuations in the causation of the seasonal variation, within the small number of years since his measurements had begun. However it is now understood that allowance should be made for about 50% absorption by the oceanic sinks.^[17] Keeling father and son have contributed to the work which has since yielded a much fuller understanding of the relation between atmospheric CO₂ and the entirety of the carbon cycle, including the effects of land and ocean sinks.^[18]

On May 9, 2013, the daily mean concentration of carbon dioxide in the atmosphere measured at Mauna Loa surpassed 400 parts per million (ppm) for the first time since measurements began in 1958.^[19]

2 See also

- Carbon dioxide in Earth's atmosphere
- Global Warming

3 References

[1] Briggs, Helen (December 1, 2007). "50 years on: The Keeling Curve legacy". *BBC News*.

- [2] Rose Kahele (October–November 2007). "Behind the Inconvenient Truth". *Hana Hou!* vol. 10, No. 5.
- [3] The Early Keeling Curve. Scripps CO2 Program
- [4] C. D. Keeling, *The Concentration and Isotopic Abundances of Carbon Dioxide in the Atmosphere*, Tellus, 12, 200-203, 1960
- [5] Keeling, Charles D. (1998). "Rewards and Penalties of Monitoring the Earth". *Annual Review of Energy and the Environment* 23: 25–82. doi:10.1146/annurev.energy.23.1.25.
- [6] ESRL Web Team. "ESRL Global Monitoring Division - Global Greenhouse Gas Reference Network". Esrl.noaa.gov. Retrieved 2014-05-26.
- [7] Trends in Atmospheric Carbon Dioxide - Mauna Loa. National Oceanic & Atmospheric Administration.
- [8] Globally averaged marine surface monthly mean data. National Oceanic & Atmospheric Administration.
- [9] Neftel, A.; Moor, E.; Oeschger, H.; Stauffer, B. (1985). "Evidence from polar ice cores for the increase in atmospheric CO₂ in the past two centuries". *Nature* 315 (6014): 45–47. Bibcode:1985Natur.315...45N. doi:10.1038/315045a0.
- [10] Keeling, Charles D. (1978). "The Influence of Mauna Loa Observatory on the Development of Atmospheric CO₂ Research". In *Mauna Loa Observatory: A 20th Anniversary Report*. (National Oceanic and Atmospheric Administration Special Report, September 1978), edited by John Miller, pp. 36-54. Boulder, CO: NOAA Environmental Research Laboratories.
- [11] Global Stations CO₂ Concentration Trends. Scripps CO2 Program.
- [12] C.D. Keeling and T.P. Whorf (October 2004). "Atmospheric CO₂ from Continuous Air Samples at Mauna Loa Observatory, Hawaii, U.S.A.". *Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory*.
- [13] Keeling Curve, 2002, University of California, San Diego
- [14] "Sampling the Air". *The New York Times*. December 22, 2010.
- [15] "CO₂ Measurements".
- [16] Manier, Jeremy (March 30, 2008). "Researcher's work, at 50, still points to 'inconvenient truth'". *Chicago Tribune*.
- [17] Nicolas Gruber and others. "Oceanic sources and sinks of atmospheric CO₂". NASA/Journal of Geophysical Research. Retrieved 2012-03-04.
- [18] I C Prentice and others. "The Carbon Cycle and Atmospheric Carbon Dioxide". IPCC. Retrieved 2012-03-04.
- [19] "Carbon Dioxide at NOAA's Mauna Loa Observatory reaches new milestone".

4 External links

- Official Keeling Curve website. Scripps Institution of Oceanography, UC San Diego
- CO2 Now: Annually-updated version of the Keeling curve
- Climate Change Is Clear Atop Mauna Loa, NPR, Day to Day, May 1, 2007
- Scripps Institution of Oceanography CO₂-Program: Home of the Keeling Curve
- Keeling Curve throughout the years

5 Text and image sources, contributors, and licenses

5.1 Text

- **Keeling Curve** *Source:* <http://en.wikipedia.org/wiki/Keeling%20Curve?oldid=641440295> *Contributors:* William M. Connolley, Jill, Dragons flight, Muxxa, Alan Liefing, JamesMLane, Scottveirs, Geni, Piotrus, Kaldari, Urhixidur, Rich Farmbrough, Vsmith, Bender235, Danielbutler, Aranel, RJFJR, Woohookitty, GregorB, Rjwilmsi, NeonMerlin, Jimbo Donal Wales, Sperxios, Bdelisle, Splette, Bergsten, Jamesg, AdiJapan, Arthur Rubin, TechBear, SmackBot, Nil Einne, Nils Simon, Monoape, Rock nj, Bobamnertiopsis, IanOfNorwich, DUDen, Cydebot, Daven200520, Thijs!bot, Nick Number, Dimawik, Ldecola, The Sanity Inspector, RockMFR, Athaenara, Bailo26, Jorfer, DorganBot, Kevinkor2, Yorambauman, Avrau, Scrippsnews, Showman60, Sjoffutt, Mrfebruary, Drmies, Boleyn, Nathan Johnson, Thomas bonasera, Addbot, Lightbot, Zorrobot, Yobot, Heqwm2, Abdallahdjabi, Bigweeboy, Recognizance, Citation bot 1, Tom.Reding, Zo8shong, Wikielwikingo, Simplex Iswrhs, RjwilmsiBot, EmausBot, Nikhiljha, Brandmeister, Mjbmrbot, ClueBot NG, O.Koslowski, Esnascosta, Bibcode Bot, BattyBot, Dexbot, JamesonRichards, Wmatysti, Lizia7, Krotera, Monkbot and Anonymous: 44

5.2 Images

- **File:Commons-logo.svg** *Source:* <http://upload.wikimedia.org/wikipedia/en/4/4a/Commons-logo.svg> *License:* ? *Contributors:* ? *Original artist:* ?
- **File:Mauna_Loa_Observatory_from_air.jpg** *Source:* http://upload.wikimedia.org/wikipedia/commons/c/c7/Mauna_Loa_Observatory_from_air.jpg *License:* Public domain *Contributors:* <http://www.mlo.noaa.gov/aboutus/siteInformation/mlosite.html> *Original artist:* National Oceanic and Atmospheric Administration
- **File:The_Keeling_Curve.png** *Source:* http://upload.wikimedia.org/wikipedia/commons/d/d3/The_Keeling_Curve.png *License:* CC BY-SA 4.0 *Contributors:* Own work *Original artist:* Scrippsnews

5.3 Content license

- Creative Commons Attribution-Share Alike 3.0