

Casimir forces between solids can be repulsive

A decades-old prediction—that materials with certain combinations of optical properties can reverse a quantum force's direction—has been experimentally verified.

The Casimir force, a result of zero-point electromagnetic fluctuations, is a quantum mechanical effect that can influence the mesoscopic and macroscopic worlds. It acts between any two objects, but it's so weak that it has typically been considered little more than a theoretical curiosity. Usually, the interaction is attractive. But now, Harvard University's Federico Capasso and his recently graduated student Jeremy Munday have observed a repulsive Casimir force—even weaker than the attractive version and with possible applications to nanoscale technology.¹

The Casimir force is a close cousin of the van der Waals force between non-polar molecules or larger objects. When the ever-moving charges in one object create a momentary electric dipole, they can induce a dipole in a nearby object such that the two dipoles attract each other. In the van der Waals limit, the objects are close enough that the electric field propagates between them much faster than the charges can oscillate in either object. But in the Casimir regime, the objects are farther apart, the finite speed of light becomes important, and the dependence of the force on the objects' separation is suitably modified.

Another way to look at the interaction is in terms of the zero-point energy of electromagnetic waves. In infinite free space, waves can have any frequency, but between two conducting plates, they're limited to the frequencies that allow an integral number of half wavelengths to fit in the gap between the surfaces. The vacuum energy therefore depends on the plates' positions, so the plates experience a force that turns out to be attractive.

Hendrik Casimir's 1940s calculations (described by Steve Lamoreaux in *PHYSICS TODAY*, February 2007, page 40) assumed that the plates were perfect conductors.² Later on, Evgeny Lifshitz and colleagues generalized Casimir's work to real materials—nonperfect conductors and dielectrics—and considered the case in which the plates were separated by a fluid, not a vacuum.³ They found that repulsive forces should result from materials with a certain relationship among their frequency-dependent dielectric permittivities. When plates of a high-permittivity material and a low-permittivity material




Figure 1. Measuring the Casimir-Lifshitz force between a 40- μm sphere and a surface. The sphere is attached to an atomic force microscope cantilever, which is lowered toward the surface by a piezoelectric column. The position of the cantilever is monitored with a superluminescent diode (SLD) and a split-quadrant photodetector. If there is an attraction between the sphere and the plate, more of the SLD intensity is detected in quadrants C and D and less in quadrants A and B. (Adapted from ref. 4.)

are separated by a fluid of intermediate permittivity, a larger momentary polarization is induced in the fluid than in the low-permittivity plate. The fluid is drawn to the high-permittivity plate more strongly than the two plates are drawn to each other, so there's a net repulsion between the plates. Notes Capasso, "That's not a rigorous explanation, but it's one way to understand the phenomenon."

Finding repulsion

Repulsive cases of the Casimir-Lifshitz force, as it's now often called, are not uncommon. They can be found whenever a liquid completely wets a solid surface: The solid and the air or ambient vapor effectively repel each other, and the liquid spreads out to fill the gap. In a similar but more spectacular example, Casimir-Lifshitz repulsion is also responsible for liquid helium's tendency to climb the walls of its container.

Combinations of materials that produce repulsive Casimir-Lifshitz forces between solid objects are harder to come by, but they're not unknown.⁴ Indeed, some have been tested already, such as gold and Teflon separated by cyclohexane, and repulsive forces have been observed.⁵ But those forces were detected only at separations of a few nanometers or less: within the van der Waals limit, which in the gold-cyclohexane-Teflon case is determined by the plasma frequency of gold. And at such small length scales, the force can be

complicated by surface roughness and the intervening liquid's inhomogeneity—the arrangement and orientation of its individual molecules.

Capasso and Munday chose gold and silica as their solid materials. With

Bellows-Sealed Linear Translator (BLT)



Operating Instructions:

1. 
2. 
3. Repeat if necessary.

McAllister Technical Services

Manufacturers of surface analytical instruments and devices

Ph. + 208-772-9527
800-445-3688

www.mcallister.com




Figure 2. Casimir-Lifshitz forces as monitored by the cantilever deflection (in arbitrary units) as a function of the piezo displacement. A negative deflection means that the sphere was attracted toward the surface, and a positive deflection means that it was repelled. **(a)** A gold sphere and a gold surface attract each other. **(b)** A gold sphere and a

silica surface, submerged in bromobenzene, repel each other. The diagonal straight lines at the left of both plots indicate that the sphere is in contact with the surface. (Adapted from ref. 1.)

the help of Adrian Parsegian (of the National Institutes of Health in Bethesda, Maryland), an expert on the chemical physics of van der Waals forces, they eliminated or accounted for many subtle effects that could interfere with their force measurement—and that might have been what prevented previous researchers from observing repulsive forces at a longer range. For example, any static charge trapped on the silica surface would induce an image charge in the gold and thereby produce an attraction. So Capasso and Munday checked that their ultrasonic cleaning procedure left no measurable surface charge on the silica.

Using a standard method for measuring Casimir forces, as shown in figure 1, the researchers attached a 40- μm gold-coated sphere to an atomic force microscope cantilever. They then lowered the sphere over a surface of gold (for comparison) or silica and monitored any additional deflection of the cantilever using a split-quadrant photodetector and a light beam from a superluminescent diode. (The sphere-plane configuration is a bit more complicated to analyze theoretically than Casimir's original plane-plane conception, but it makes the experiment much easier, because keeping the planes exactly parallel is not a concern.) They first tried the experiment in ethanol, but ultimately changed to bromobenzene, a less polar substance that also satisfies the necessary permittivity relationship over a wide range of frequencies.

The results are shown in figure 2. As

expected, for the gold-gold configuration in figure 2a, the interaction was attractive: As the cantilever was lowered, the sphere was pulled toward the surface until it made contact, as indicated by the abrupt change in slope of the deflection curve. When the cantilever was retracted, the sphere stuck to the surface as the cantilever was raised an additional 10 nm—the attraction is stronger when the sphere and the surface are already in contact than when they're not—and then was pulled loose.

For the gold-silica combination in figure 2b, the cantilever was deflected upward during both the approach and the retraction. To convert the cantilever deflection data into a force measurement, the researchers had to isolate the effect of the Casimir-Lifshitz force from that of the hydrodynamic force of the bromobenzene on the gold sphere. Because that drag force is proportional to the velocity of the sphere, its contribution could be determined by repeating the experiment at different speeds. With the drag force thus canceled out, Capasso and Munday were able to measure repulsive forces of tens of piconewtons for sphere-surface separations of up to 40 nm.

Levitation applications

From the beginning, Capasso's interest in the Casimir-Lifshitz force was in part motivated by its relevance to microelectromechanical systems, or MEMS. If two parts of one of those tiny devices get too close to each other, they

Goodfellow
Metals & Materials



**STRAIGHT UP
OR WITH
A TWIST**



**Goodfellow has
the flexibility to deliver
just what you need.**

For *standard* catalog items
goodfellow.com

For *custom* items and
special sizes and quantities
info@goodfellow.com

To speak to a real, live person
1-800-821-2870

Small quantities FAST!

© 2008 Goodfellow Corporation

stick together and render the system inoperable. The phenomenon is called stiction, and it's thought that the Casimir–Lifshitz force is responsible. Reducing or eliminating the attractive Casimir–Lifshitz force among MEMS components could alleviate the problem of stiction as MEMS are further miniaturized. And actually reversing the force into a repulsion allows the intriguing possibility of levitating one object above another. If a levitation gap can be achieved that's larger than any surface roughness features (typically

around 15 nm), it could allow the creation of very low-friction force sensors, bearings, or other devices.

Capasso also hopes that Casimir levitation will help him observe a quantum electrodynamic torque between optically anisotropic materials. By levitating one disk of birefringent material above another, he says, it should be possible to rotate the top disk with circularly polarized light and see that the disks' principal optical axes tend to align when the light is removed.⁴

Johanna Miller

References

1. J. N. Munday, F. Capasso, V. A. Parsegian, *Nature* **457**, 170 (2009).
2. H. B. G. Casimir, D. Polder, *Phys. Rev.* **73**, 360 (1948); H. B. G. Casimir, *J. Chim. Phys.* **46**, 407 (1949); H. B. G. Casimir, *Proc. K. Ned. Akad. Wet.* **51**, 793 (1948).
3. E. M. Lifshitz, *J. Exp. Theor. Phys.* **2**, 73 (1956); I. E. Dzyaloshinskii, E. M. Lifshitz, L. P. Pitaevskii, *Adv. Phys.* **10**, 165 (1961).
4. F. Capasso, J. N. Munday, D. Iannuzzi, H. B. Chan, *IEEE J. Select. Top. Quant. Electron.* **13**, 400 (2007).
5. A. A. Feiler, L. Bergstrom, M. W. Rutland, *Langmuir* **24**, 2274 (2008).

physics update

These items, with supplementary material, first appeared at <http://www.physicstoday.org>.


Ice acoustics for detecting neutrinos. Several experiments are operating or being built to detect astrophysical neutrinos. Ranging up to about a cubic kilometer in size, those experiments are embedded in ice or in a liquid such as water, where they watch for telltale flashes of Cherenkov radiation. (See the article by Francis Halzen and Spencer Klein in *PHYSICS TODAY*, May 2008, page 29.) But the highest-energy neutrinos, with energies of an exaelectron volt (1EeV = 10^{18} eV) or higher, are so scarce that



installations spanning 100 km³, along with massive numbers of expensive photomultiplier tubes, would be needed to collect adequate event statistics in a reasonable time. So other detection schemes are being explored, one of which involves acoustics: When a very-high-energy neutrino interacts with water or ice, a sudden localized thermal expansion occurs and the resulting wave propagates farther than the light flashes. To explore that method, the Aachen Acoustic Laboratory was set up in late 2007 and its first experiment made a precise measurement of the speed of sound in ice that is entirely devoid of bubbles and cracks. The Aachen physicists carefully positioned an array of sensors—six detectors and one emitter—in a 3-m³ water tank (shown here) equipped with a freeze-control unit and a degassing system. The difference in arrival times of an acoustic pulse at adjacent receivers determined the speed of sound. Between 0 °C and –17 °C, where they took measurements, the speed ranged from about 3840 m/s to 3890 m/s, agreeing well with earlier laboratory experiments. The team is also part of SPATS (the South Pole Acoustic Test Setup), which is currently obtaining complementary in situ measurements. (C. Vogt, K. Laihem, C. Wiebusch, *J. Acoust. Soc. Am.* **124**, 3613, 2008.)

—SGB

Tuning vibrations for label-free biological imaging. To map molecules in cells and tissue, researchers prefer biomedical imaging techniques that rely solely on the intrinsic responses of chemical bonds to optical stimulation. Although fluorescence microscopy and other chemical tagging methods yield high-resolution images, they also introduce foreign species or syn-



thetic derivatives that can alter the dynamics of intracellular processes. Spontaneous Raman scattering, which uses a single laser beam to excite the vibrational and rotational modes in chemical bonds, requires no chemical labels but generates a weak signal that gets muddled by Rayleigh scattering. A more sensitive technique known as coherent anti-Stokes Raman scattering uses multiple laser beams to generate coherent optical signals that enhance resonant frequencies in the sample; that method, however, also produces nonresonant background noise. Recently a team led by Harvard University chemist Sunney Xie demonstrated a new technique based on stimulated Raman scattering that tunes the difference between the frequencies of two laser beams to match a desired molecule's resonant frequency, thus amplifying the Raman signal. The measurable intensities of the transmitted beams change only when a match occurs; nonresonant signals are not picked up. The images show the top view (a) and the depth profile (b) of an acne medication (blue) that penetrated a mouse's skin, thus demonstrating the potential of the new technique to monitor drug delivery. (C. W. Freudiger et al., *Science* **322**, 1857, 2008.)

—JNAM

Heating the Sun's corona. It's one of the great natural mysteries: How do the Sun's corona and wind become thousands of times hotter than the Sun's surface? Somehow, energy makes its way up into the corona against a temperature gradient and is converted to heat. A new analysis of data collected by NASA's *Wind* spacecraft doesn't solve the mystery, but it is consistent with a popular explanation. The analysis was done by Justin Kasper of the Harvard-Smithsonian Center for Astrophysics, Alan Lazarus of MIT, and Peter Gary of Los Alamos National Laboratory. They looked at 14 years of in situ observations of particles and fields made as *Wind* flew in and out of the solar wind. The team focused on the two most abundant ion species in the solar wind, H⁺ and He²⁺. Because He²⁺ is four times heavier than H⁺ and carries twice the charge, the two species' kinematics can discriminate among various models for transport and heating. Kasper, Lazarus, and Gary found strong evidence for one picture of coronal heating: Ions are carried upward by magnetohydrodynamic